

Supporting Information for:

Tumor-Derived Extracellular Vesicles Breach the Intact Blood Brain Barrier *via* Transcytosis

Golnaz Morad^{†,‡}, Christopher V. Carman[§], Elliott J. Hagedorn^{¶,††}, Julie R. Perlin^{¶,††}, Leonard I. Zon^{¶,††}, Nur Mustafaoglu^{‡‡}, Tae-Eun Park^{‡‡,§§}, Donald E. Ingber^{†,‡‡,¶¶}, Cassandra C. Daisy[†], Marsha A. Moses^{*,†,‡,†††}

[†] Vascular Biology Program, Boston Children's Hospital, Boston, MA, 02115 USA;

[‡] Department of Surgery, Harvard Medical School, Boston, MA, 02115 USA;

[§] Molecular and Integrative Physiological Sciences Program, Harvard T.H. Chan School of Public Health, Boston, MA, 02115 USA;

[¶] Stem Cell Program and Division of Hematology/Oncology, Boston Children's Hospital and Dana Farber Cancer Institute, Howard Hughes Medical Institute, Harvard Stem Cell Institute, Harvard Medical School, Boston, MA, 02115 USA;

^{††} Department of Stem Cell and Regenerative Biology and Harvard Stem Cell Institute, Harvard University, Cambridge, MA, 02115 USA;

^{‡‡} Wyss Institute for Biologically Inspired Engineering at Harvard University, Boston, MA, 02115 USA;

^{§§} Ulsan National Institute of Science and Technology, Ulsan, South Korea;

^{¶¶} Harvard John A. Paulson School of Engineering and Applied Sciences, Cambridge, MA, 02138 USA;

^{†††} Department of Surgery, Boston Children's Hospital, Boston, MA, 02115 USA.

*To whom correspondence should be addressed:

Marsha A. Moses, Ph.D.

Vascular Biology Program

Boston Children's Hospital

The Karp Family Research Building


300 Longwood Avenue, Boston MA 02115

Phone (617) 919-2207


Fax (617) 730-0231

Email: marsha.moses@childrens.harvard.edu


Supplementary Figures



Supplementary Figure S1. Characterization of the cell lines and the isolated EVs. (a) *in vivo* luminescent imaging of metastases following intracardiac injection of parental and brain-seeking MDA-MB-231 cells. (b) Nanoparticle tracking analysis of the size of P-EVs and Br-EVs. (c) Representative western blot images of EV markers CD9, CD63, Alix, and the golgi marker, GM130. (d) The percentage of the mice that developed brain metastases following treatment with PBS, P-EVs, and Br-EVs (n=7 mice per group).



Supplementary Figure S2. Characterization of the *in vitro* BBB model and the transcytosed EVs. (a) Schematic showing static BBB model preparation and transcytosis experiments. (b) Representative images of brain endothelial cells immunostained with anti-ZO-1 antibody following treatment with cAMP and Ro 20-1724 (3 independent experiments). (c) Fold change in permeability coefficient of brain endothelial monolayer to 10 KDa (upper graph) and 70 KDa (lower graph) dextran following treatment with cAMP and Ro 20-1724 (mean \pm SD; 3 independent experiments). Statistical analysis was performed using one-way ANOVA with Tukey's correction for multiple comparisons. (d) Luminescent intensity of in density fractions following density gradient fractionation of luciferase-labeled Br-EVs. (e) Electron microscopy images and (F) quantification of the size of EVs isolated from the low density (15% Optiprep) and high density (25% Optiprep) fractions. Statistical analysis was performed using Student's t-test. In all panels, ns, not significant; * $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$.



Supplementary Figure S3. Colocalization of Br-EVs with caveolin, eea1, and TGN46. (a) Representative fluorescence microscopy image of brain endothelial cells immunostained with anti-caveolin 1 antibody from 3 independent experiments. Scale bar, 25 μm . (b) Representative fluorescence microscopy image of brain endothelial cells immunostained with anti-eea1 antibody from 3 independent experiments. Right panels show the magnification of the area in the white square. Scale bar, 25 μm . (c) Representative fluorescence microscopy image of brain endothelial cells immunostained with anti-TGN46 antibody from 3 independent experiments. Scale bar, 25 μm .